



11030
BB R1500 ALL PURPOSE

MEDICATED

Feed to mature reproducing beef cows: For improved feed efficiency; for prevention and control of coccidiosis due to E. bovis and E. zuernii.

ACTIVE DRUG INGREDIENT

Monensin (as Monensin sodium) 1512.0 g/ton

Mineral for Beef Cattle.

GUARANTEED ANALYSIS

| | | | |
|----------------|-----------|---------|-------|
| Calcium (Ca) | (Minimum) | 18.6 | % |
| Calcium (Ca) | (Maximum) | 23.6 | % |
| Phosphorus (P) | (Minimum) | 0.10 | % |
| Salt (NaCl) | (Minimum) | 22.6 | % |
| Salt (NaCl) | (Maximum) | 27.6 | % |
| Magnesium (Mg) | (Minimum) | 1.5 | % |
| Potassium (K) | (Minimum) | 0.2 | % |
| Copper (Cu) | (Minimum) | 501.1 | ppm |
| Iodine | (Minimum) | 59.4 | ppm |
| Selenium (Se) | (Minimum) | 20.0 | ppm |
| Zinc (Zn) | (Minimum) | 1,810.3 | ppm |
| Manganese (Mn) | (Minimum) | 1,114.1 | ppm |
| Vitamin A | (Minimum) | 222,211 | IU/Lb |
| Vitamin E | (Minimum) | 152 | IU/Lb |

INGREDIENTS

Calcium Carbonate, Salt, Processed Grain By-Products, Magnesium Oxide, Midwest Ag Beef Trace Min., Vegetable Fat, Thiamine Mononitrate, Vitamin A Supplement, Roughage Products, Vitamin E Supplement, Vitamin D3 Supplement, Mineral Oil, Sodium Selenite, Natural & Artificial Flavor.

WARNING: A withdrawal time has not been established for preruminating calves. Do not use in calves to be processed for veal. **CAUTION:** Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended by the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

LIMITATIONS: Feed to mature reproducing beef cows. Feed as supplemental feed, either hand-fed in a minimum of 1 pound of feed or mixed in a total ration. For improved feed efficiency, feed continuously at a rate of 50 to 200 milligrams Monensin per head per day. For prevention and control of coccidiosis, feed at a rate of 0.14 to 0.42 milligram per pound of body weight per day, depending upon severity of challenge, up to a maximum of 200 milligrams per head per day. During first 5 days of feeding, cattle should receive no more than 100 milligrams per head per day.

DIRECTIONS FOR USE

Feed as directed by Countryside Feed, LLC. Feed at a rate of .1 lb to .48 lb per head per day to provide between 75 and 360 mg of Monensin daily.



Manufactured By:
Countryside Feed, LLC.
Seneca, KS 66538

Bulk - or - 50 lb (22.68 kg) Net Weight

MEDICATED