



60153
BB MGA RATION- 1/2#/HEAD

MEDICATED

For suppression of estrus (heat) in heifers intended for reproduction.

ACTIVE DRUG INGREDIENTS

Melengestrol Acetate 0.00022 %

For Beef Replacement Heifer Cattle on Pasture.

GUARANTEED ANALYSIS

Crude Protein	(Minimum)	7.5	%
Crude Fat	(Minimum)	3.6	%
Crude Fiber	(Maximum)	2.6	%
Calcium (Ca)	(Minimum)	0.02	%
Calcium (Ca)	(Maximum)	0.02	%
Phosphorus (P)	(Minimum)	0.3	%
Potassium (K)	(Minimum)	0.3	%

INGREDIENTS

Grain Products.

CAUTION: Withdrawal periods of 3 to 5 days should be avoided to prevent the possibility that heifers may come into estrus (heat) at loading time. Not effective in steers or spayed heifers.

Heifers intended for reproduction; do not exceed 24 days of feeding. A reduced conception rate can be expected if heifers are bred at estruses observed within 1 to 12 days after withdrawal of Melengestrol Acetate, whereas heifers bred at subsequent observed estrus are expected to have normal conception rates.

LIQUID LIMITATIONS: The liquid Type B feed and liquid Type C feed must have a pH of 4.0 to 8.0 and must be agitated or re-circulated thoroughly before using to insure proper mixing of ingredients. Liquid Melengestrol Acetate may not be mixed with Oxytetracycline in a common liquid feed supplement.

DIRECTIONS FOR USE

Feed as directed by Countryside Feed, LLC. Feed .5 lbs per head per day which provides .5 mg of MGA.



Manufactured By:
Countryside Feed, LLC.
Seneca, KS 66538

Bulk - or - 50 lb (22.68 kg) Net Weight

MEDICATED