



68

BB COW SUPP 25%

MEDICATED

Growing cattle on pasture or in dry lot (stocker and feeder cattle and dairy and beef replacement heifers): For increased rate of weight gain. For prevention and control of coccidiosis caused by Eimeria bovis and Eimeria zuernii.

ACTIVE DRUG INGREDIENT

Monensin (as Monensin sodium) 362.8 g/ton

For Beef Cattle on Pasture.

GUARANTEED ANALYSIS

| | Crude Protein* | (Minimum) | 25.1 | % |
|--|----------------|-----------|--------|-------|
| * This includes not more than 9.8 base % equivalent protein from non-protein nitrogen. | | | | |
| | Crude Fat | (Minimum) | 3.7 | % |
| | Crude Fiber | (Maximum) | 7.7 | % |
| | Calcium (Ca) | (Minimum) | 3.1 | % |
| | Calcium (Ca) | (Maximum) | 4.1 | % |
| | Phosphorus (P) | (Minimum) | 1.0 | % |
| | Salt (NaCl) | (Minimum) | 20.4 | % |
| | Salt (NaCl) | (Maximum) | 25.4 | % |
| | Potassium (K) | (Minimum) | 0.6 | % |
| | Vitamin A | (Minimum) | 40,000 | IU/Lb |

INGREDIENTS

Processed Grain By-Products, Salt, Roughage Products, Calcium Chloride, UREA, Monocalcium Phosphate, Plant Protein Products, Calcium Carbonate, Magnesium Oxide, Midwest Ag Beef Trace Min., Vitamin A Supplement, Vitamin E Supplement, Vitamin D3 Supplement, Mineral Oil, Sodium Selenite.

CAUTION: A withdrawal time has not been established for preruminating calves. Do not use in calves to be processed for veal. Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

LIMITATIONS: For increased rate of weight gain, feed at a rate of 50 to 200 milligrams Monensin per head per day in not less than 1 pound of feed or, after the 5th day, feed at a rate of 400 milligrams per head per day every other day in not less than 2 pounds of feed. For prevention and control of coccidiosis, feed at a rate of 0.14 to 0.42 milligram per pound of body weight per day, depending on severity of challenge, up to 200 milligrams per head per day. During first 5 days of feeding, cattle should receive no more than 100 milligrams per day in not less than 1 pound of feed.

CAUTION: USE AS DIRECTED.

DIRECTIONS FOR USE

Feed as directed by Countryside Feed, LLC. Feed 1 lb per head per day which provides 180 mg of Monensin.



Manufactured By:
Countryside Feed, LLC.
Seneca, KS 66538

Bulk - or - 50 lb (22.68 kg) Net Weight

MEDICATED