



70574
4-H Show Pig Grower Denagard

MEDICATED

For control of swine dysentery associated with Brachyspira (formerly Serpulina or Treponema) hyodysenteriae susceptible to tiamulin. Prevents the development of house flies, stable flies, face flies, and horn flies in the manure of treated swine.

ACTIVE DRUG INGREDIENT	
Tiamulin hyd	35 g/ton
ACTIVE INGREDIENT	
Diflubenzuron	7.9 mg/lb

For Grower Pigs.

GUARANTEED ANALYSIS

Crude Protein	(Minimum)	16.5	%
Lysine	(Minimum)	1.09	%
Methionine	(Minimum)	0.3	%
Threonine	(Minimum)	0.8	%
Crude Fat	(Minimum)	5.0	%
Crude Fiber	(Maximum)	3.7	%
Calcium (Ca)	(Minimum)	0.70	%
Calcium (Ca)	(Maximum)	1.20	%
Phosphorus (P)	(Minimum)	0.56	%
Salt (NaCl)	(Minimum)	0.5	%
Salt (NaCl)	(Maximum)	0.9	%
Magnesium (Mg)	(Minimum)	0.2	%
Potassium (K)	(Minimum)	0.74	%
Copper (Cu)	(Minimum)	209.7	ppm
Selenium (Se)	(Minimum)	0.4	ppm
Zinc (Zn)	(Minimum)	219.2	ppm
Iron (Fe)	(Minimum)	179.4	ppm
Manganese (Mn)	(Minimum)	19.3	ppm
Vitamin A	(Minimum)	5,589	IU/Lb
Vitamin D3	(Minimum)	800	IU/Lb
Vitamin E	(Minimum)	37	IU/Lb
Biotin	(Minimum)	61.6	Mcg/Lb
Riboflavin	(Minimum)	4.9	Mg/Lb
Niacin	(Minimum)	30.6	Mg/Lb
Vitamin B12	(Minimum)	15	Mcg/Lb
Panto Acid	(Minimum)	13.7	Mg/Lb
Phytase	(Minimum)	3	FTU/lb

INGREDIENTS

Grain Products, Plant Protein Products, Animal Protein Products, Roughage Products, Animal Fat, Calcium Carbonate, Monocalcium Phosphate, Salt, Choline Chloride, Vitamin E Supplement, Niacin Supplement, d-Calcium Pantothenate, Vitamin A Acetate, Folic Acid, d-Biotin, Riboflavin Supplement, Menadione Sodium Bisulfite Complex, Pyridoxine Hydrochloride, Vitamin D3 Supplement, Vitamin B12 Supplement, Thiamine Mononitrate, Ferrrous Sulfate, Zinc Oxide, Manganous Oxide, Selenium Yeast, Copper Sulfate, Dried Aspergillus niger Fermentation Product, Mineral Oil, Dried Schizosaccharomyces pombe Fermentation Product, Sodium Selenite and Ethylenediamine Dihydroiodide, L-Lysine, Calcium Propionate, L-Threonine, DL-Methionine, _

WARNING: Withdraw 2 days before slaughter.

LIMITATIONS: Feed continuously as sole ration on premises with a history of swine dysentery but where signs of disease have not yet occurred or following approved treatment of disease.

SPECIAL CONSIDERATIONS: Swine being treated with Tiamulin should not have access to feeds containing polyether ionophores (e.g., Lasalocid, Monensin, Narasin, Salinomycin, or Semduramycin) as adverse reactions may occur. If signs of toxicity occur, discontinue use. The effects of Tiamulin on swine reproductive performance, pregnancy, and lactation have not been determined. Use as sole source of tiamulin.

CAUTION: Keep out of reach of children. Discard empty container according to local regulations. Never reuse empty container.

DIRECTIONS FOR USE

Feed as Sole Ration.



Manufactured By:
Countryside Feed, LLC.
Seneca, KS 66538

Bulk - or - 50 lb (22.68 kg) Net Weight

MEDICATED