



# 71116 Pork Builder Jump Start Denagard

#### **MEDICATED**

For control of swine dysentery associated with Brachyspira (formerly Serpulina or Treponema)hyodysenteriae susceptible to tiamulin. Prevents the development of house flies, stable flies, face flies, and horn flies in the manure of treated swine.

### 

## For Grower Pigs.

#### **GUARANTEED ANALYSIS** Crude Protein (Minimum) 192 % Lysine (Minimum) 1.36 % Crude Fat (Minimum) 4.7 % Crude Fiber (Maximum) 3.8 % Calcium (Ca) (Minimum) 0.69 % Calcium (Ca) (Maximum) 1.19 % Phosphorus (P) % (Minimum) 0.64 Salt (NaCI) (Minimum) 0.6 % Salt (NaCI) (Maximum) 1.0 % Zinc (Zn) (Minimum) 221.4 ppm Vitamin A (Minimum) 5,506 IU/Lb Vitamin F (Minimum) 36 IU/Lb **INGREDIENTS**

Grain Products, Plant Protein Products, Animal Protein Products, Roughage Products, Animal Fat, Monocalcium Phosphate, Calcium Carbonate, Salt, L-Lysine, Choline Chloride, Vitamin E Supplement, Niacin Supplement, d-Calcium Pantothenate, Vitamin A Acetate, Folic Acid, d-Biotin, Riboflavin Supplement, Menadione Sodium Bisulfite Complex, Pyridoxine Hydrochloride, Vitamin D3 Supplement, Vitamin B12 Supplement, Thiamine Mononitrate, Ferrous Sulfate, Zinc Oxide, Manganous Oxide, Selenium Yeast, Copper Sulfate, Dried Aspergillus niger Fermentation Product, Mineral Oil, Dried Schizosaccharomyces pombe Fermentation Product, Sodium Selenite and Ethylenediamine Dihydroiodide., DL-Methionine, L-Threonine, \_.

WARNING: Withdraw 2 days before slaughter.

**LIMITATIONS:** Feed continuously as sole ration on premises with a history of swine dysentery but where signs of disease have not yet occurred or following approved treatment of disease. **SPECIAL CONSIDERATIONS:** Swine being treated with Tiamulin should not have access to feeds containing polyether ionophores (e.g., Lasalocid, Monensin, Narasin, Salinomycin, or Semduramycin) as adverse reactions may occur. If signs of toxicity occur, discontinue use. The effects of Tiamulin on swine reproductive performance, pregnancy, and lactation have not been determined. Use as sole source of tiamulin.

**CAUTION:** Keep out of reach of children. Discard empty container according to local regulations. Never reuse empty container.

DIRECTIONS FOR USE Feed as Sole Ration.



Bulk - or - 50 lb (22.68 kg) Net Weight