



80140
 MWAS 0-40 Day Lamb Ration

MEDICATED

For the prevention of coccidiosis in young sheep caused by Eimeria ovinoidalis, E. crandallis, E. parva and E. bakuensis.

ACTIVE DRUG INGREDIENT

Decoquinatate 14 g/ton

Starter Feed for Goat or Sheep.

GUARANTEED ANALYSIS

	Crude Protein*	(Minimum)	19.5	%
* This includes not more than 0.9 base % equivalent protein from non-protein nitrogen.				
	Crude Fat	(Minimum)	2.5	%
	Crude Fiber	(Maximum)	2.6	%
	Calcium (Ca)	(Minimum)	0.68	%
	Calcium (Ca)	(Maximum)	1.18	%
	Phosphorus (P)	(Minimum)	0.36	%
	Salt (NaCl)	(Minimum)	1.6	%
	Salt (NaCl)	(Maximum)	2.1	%
	Vitamin A	(Minimum)	2,083	IU/Lb
	Vitamin E	(Minimum)	14	IU/Lb

INGREDIENTS

Grain Products, Plant Protein Products, Molasses Products, Calcium Carbonate, Salt, Ammonium Chloride.

WARNING: Do not feed to sheep producing milk for food.

CAUTION: Do not use in feeds containing bentonite.

LIMITATIONS: Feed at least 28 days during periods of exposure to coccidiosis or when experience indicates that it is likely to be a hazard. Feed Type C feed or milk replacer at a rate to provide 22.7 mg decoquinatate per 100 lb (0.5 mg/kg) bodyweight. Type C feed may be prepared from dry or liquid type B feed.

CAUTION: USE AS DIRECTED.

DIRECTIONS FOR USE

Feed as directed by Countryside Feed, LLC. Feed 2 lbs per head per day which provides 27.2 mg of Deccox.



Manufactured By:
 Countryside Feed, LLC.
 Seneca, KS 66538

Bulk - or - 50 lb (22.68 kg) Net Weight

MEDICATED