



80144  
 MWAS 41-80 day Lamb Grower

MEDICATED

For the prevention of coccidiosis in young sheep caused by Eimeria ovinoidalis, E. crandallis, E. parva and E. bakuensis.

ACTIVE DRUG INGREDIENT

Decoquinatate ..... 14 g/ton

Grower Feed for Goat and Sheep.

GUARANTEED ANALYSIS

Crude Protein\* (Minimum) 15.7 %

\* This includes not more than 0.9 base % equivalent protein from non-protein nitrogen.

Crude Fat	(Minimum)	2.7	%
Crude Fiber	(Maximum)	2.5	%
Calcium (Ca)	(Minimum)	0.64	%
Calcium (Ca)	(Maximum)	0.89	%
Phosphorus (P)	(Minimum)	0.33	%
Salt (NaCl)	(Minimum)	1.6	%
Salt (NaCl)	(Maximum)	2.1	%
Vitamin A	(Minimum)	2,159	IU/Lb

INGREDIENTS

Grain Products, Plant Protein Products, Molasses Products, Calcium Carbonate, Salt, Ammonium Chloride, Thiamine Mononitrate.

WARNING: Do not feed to sheep producing milk for food.

CAUTION: Do not use in feeds containing bentonite.

LIMITATIONS: Feed at least 28 days during periods of exposure to coccidiosis or when experience indicates that it is likely to be a hazard. Feed Type C feed or milk replacer at a rate to provide 22.7 mg decoquinatate per 100 lb (0.5 mg/kg) bodyweight. Type C feed may be prepared from dry or liquid type B feed.

CAUTION: USE AS DIRECTED.

DIRECTIONS FOR USE

Feed as directed by Countryside Feed, LLC. Feed 3 lb per head per day to provide 20 mg Deccox.



Manufactured By:  
 Countryside Feed, LLC.  
 Seneca, KS 66538

Bulk - or - 50 lb (22.68 kg) Net Weight

**MEDICATED**