

14% PELLETTED HORSE FEED

FEED FOR MAINTENANCE OF HORSES

GUARANTEED ANALYSIS

Crude Protein (Min).....	14.00 %
Lysine (Min).....	0.60 %
Crude Fat (Min).....	3.00 %
Crude Fiber (Max).....	8.00 %
Acid Detergent Fiber (ADF) (Max).....	11.00 %
Neutral Detergent Fiber (NDF) (Max).....	28.00 %
Calcium (Ca) (Min).....	0.80 %
Calcium (Ca) (Max).....	1.30 %
Phosphorus (P) (Min).....	0.60 %
Magnesium (Mg) (Min).....	0.20 %
Copper (Cu) (Min).....	50.00 ppm
Selenium (Se) (Min).....	0.40 ppm
Zinc (Zn) (Min).....	150.00 ppm
Vitamin A (Min).....	5000 IU/LB
Vitamin E (Min).....	65 IU/LB
Biotin (Min).....	0.20 MG/LB

INGREDIENTS

Processed Grain By-Products, Grain Products, Forage Products, Plant Protein Products, Molasses Products, Calcium Carbonate, Soybean Oil, Lignin Sulfonate, Salt, DL-Methionine, Biotin, Vitamin E Supplement, Copper Sulfate, Zinc Oxide, Vitamin D3 Supplement, Vitamin A Supplement, Calcium Iodate, Magnesium Oxide, Cobalt Carbonate, Ferrous Carbonate, Sodium Selenite, Manganous Oxide.

35WU-STJ-G 2

DIRECTIONS

Feed with hay and/or pasture to make a complete ration for horses. Feed mature horses at the rate of 0.5 to 1 pound per 100 pounds body weight, along with 1 to 1.5 pounds of hay or equivalent pasture per 100 pounds body weight daily.

After observing your horse for a period of time, you may want to increase or decrease the daily feeding to keep your horse in the desired condition. The amount required will vary, depending on the quality and amount of hay or pasture consumed. Do not feed more than 0.5 lbs per 100 lbs body weight of this product at a single feeding.

RECOMMENDATIONS FOR FEEDING HORSES:

The recommended feeding rate should not be exceeded. Do not feed free-choice. Any feed changes should be made gradually over a period of 7 to 10 days. Changes in the rate of feeding should not exceed one pound per day for each horse. Feed at regular times, at least twice daily, with three daily feedings preferred. Reduce and/or delay feeding a horse which is hot, excited or showing pain, fever or diarrhea. Let horses feed in a natural position from troughs with large bottoms, placed at normal head

See Reverse Side For Directions for Use and Precautionary Statements

MANUFACTURED FOR

Countryside Feed, LLC

1972 KS-187, Seneca, KS 66538

Net Weight 50 lb (22.67 kg)

3006802-206

14% PELLETTED HORSE FEED

35WU

height or lower. Have plenty of fresh, clean water available at all times.

The recommended feeding rate should not be exceeded. Do not feed free-choice. Any feed changes should be made gradually over a period of 7 to 10 days. Changes in the rate of feeding should not exceed one pound per day for each horse. Feed at regular times, at least twice daily, with three daily feedings preferred.

Reduce and/or delay feeding a horse which is hot, excited or showing pain, fever or diarrhea. Let horses feed in a natural position from troughs with large bottoms, placed at normal head height or lower. Have plenty of fresh, clean water available at all times.

Maintain an effective control program for internal parasites. Be sure that horses are free of dental problems. Prevent the rapid eating by the horses of any feedstuff. Observe your horses' condition daily. Consult your veterinarian if any problems arise.

CAUTION

Store in a dry, well-ventilated area protected from rodents and insects. Do not feed moldy or insect-infested feed to animals as it may cause illness, performance loss or death.

USE ONLY AS DIRECTED