



71116

Pork Builder Jump Start Denagard

MEDICATED

For control of swine dysentery associated with *Brachyspira* (formerly *Serpulina* or *Treponema*) hyodysenteriae susceptible to tiamulin. Prevents the development of house flies, stable flies, face flies, and horn flies in the manure of treated swine.

ACTIVE DRUG INGREDIENT

Tiamulin hyd 35 g/ton

ACTIVE INGREDIENT

Diflubenzuron 6.08 mg/lb

For Grower Pigs.

GUARANTEED ANALYSIS

Crude Protein	(Minimum)	19.2	%
Lysine	(Minimum)	1.36	%
Crude Fat	(Minimum)	4.7	%
Crude Fiber	(Maximum)	3.8	%
Calcium (Ca)	(Minimum)	0.7	%
Calcium (Ca)	(Maximum)	1.2	%
Phosphorus (P)	(Minimum)	0.6	%
Salt (NaCl)	(Minimum)	0.6	%
Salt (NaCl)	(Maximum)	1.0	%
Zinc (Zn)	(Minimum)	221.4	ppm
Vitamin A	(Minimum)	5,506	IU/Lb
Vitamin E	(Minimum)	36	IU/Lb

INGREDIENTS

Grain Products, Plant Protein Products, Animal Protein Products, Animal Fat, Roughage Products, Monocalcium Phosphate, Calcium Carbonate, Salt, L-Lysine, Choline Chloride, Vitamin E Supplement, Niacin Supplement, d-Calcium Pantothenate, Vitamin A Acetate, Folic Acid, d-Biotin, Riboflavin Supplement, Menadione Sodium Bisulfite Complex, Pyridoxine Hydrochloride, Vitamin D3 Supplement, Vitamin B12 Supplement, Thiamine Mononitrate, Ferrous Sulfate, Zinc Oxide, Manganous Oxide, Selenium Yeast, Copper Sulfate, Dried *Aspergillus niger* Fermentation Product, Mineral Oil, Dried *Schizosaccharomyces pombe* Fermentation Product, Sodium Selenite and Ethylenediamine Dihydroiodide., L-Threonine, DL-Methionine, _.

WARNING: Withdraw 2 days before slaughter.

LIMITATIONS: Feed continuously as sole ration on premises with a history of swine dysentery but where signs of disease have not yet occurred or following approved treatment of disease.

SPECIAL CONSIDERATIONS: Swine being treated with Tiamulin should not have access to feeds containing polyether ionophores (e.g., Lasalocid, Monensin, Narasin, Salinomycin, or Semduramycin) as adverse reactions may occur. If signs of toxicity occur, discontinue use. The effects of Tiamulin on swine reproductive performance, pregnancy, and lactation have not been determined.

Use as sole source of tiamulin.

CAUTION: Keep out of reach of children. Discard empty container according to local regulations. Never reuse empty container.

DIRECTIONS FOR USE

Feed as Sole Ration.



Manufactured By:
Countryside Feed, LLC.
Seneca, KS 66538
71116

Bulk - or - 50 lb (22.68 kg) Net Weight

MEDICATED