



Purina® Calf Startena Coarse DQ 0.0032%

Specie: Cattle **Type of Feed:** Type C Medicated **Form of Feed:** Pellet

General Description: For the prevention of coccidiosis in ruminating and non-ruminating calves, including veal calves, and cattle caused by *Eimeria bovis* and *E. zuernii*.

PURINA® CALF STARTENA COARSE DQ 0.0032% MEDICATED

SUPPLEMENT FEED FOR DAIRY STARTER -3 DAYS TO 3 MOS IN CONFINEMENT
CAUTION: USE ONLY AS DIRECTED

ACTIVE DRUG INGREDIENTS

Decoquinatate 0.0032 %
14.5 (mg/lb)

GUARANTEED ANALYSIS

Crude Protein, (Min) 16.00 %
Crude Fat, (Min) 2.00 %
Crude Fiber, (Max) 8.00 %
Acid Detergent Fiber (ADF), (Max) 12.00 %
Calcium (Ca), (Min) 1.00 %
Calcium (Ca), (Max) 1.50 %
Phosphorus (P), (Min) 0.50 %
Selenium (Se), ppm, (Min) 0.30
Vitamin A, I.U./lb, (Min) 3,000

INGREDIENTS

Grain Products, Processed Grain By-Products, Plant Protein Products, Molasses Products, Sodium Sesquicarbonate, Calcium Carbonate, Soybean Oil, Salt, Citric Acid, Propionic Acid (a Preservative), Vitamin A Supplement, Vitamin D3 Supplement, Vitamin E Supplement, Zinc Sulfate, Manganese Sulfate, Riboflavin Supplement, Copper Sulfate, Ferrous Sulfate, Dicalcium Phosphate, Monocalcium Phosphate, Cobalt Carbonate, Vitamin B12 Supplement, Ethylenediamine Dihydroiodide, Sodium Selenite, Natural and Artificial Flavors

FEEDING DIRECTIONS

Feed at the rate of 1.56 lbs per 100 lbs of bodyweight to provide 22.7 mg Decoquinatate per 100 pounds of body weight (0.5 mg/kg) a day.

Feed at least 28 days during period of exposure to coccidiosis or when it is likely to be a hazard.

CAUTION

Bentonite should not be used in decoquinatate feeds. Store in a dry, well ventilated area protected from rodents and insects. Do not feed moldy or insect-infested feed to animals as it may cause illness, performance loss or death. This product was made in a feed manufacturing facility and does not handle or store products containing animal proteins prohibited in ruminant feed. USE ONLY AS DIRECTED.

WARNING

Do not feed to cows producing milk for food.

IMPORTANT:

This product is to be fed only to the animal species as directed on this label. Follow these management practices:

1. Provide a source of fresh clean water at all times.
2. To ensure freshness, offer only the amount of feed recommended.
3. Clean feeder pails daily.
4. Proper housing and sanitation are important to help prevent calf diarrhea and pneumonia. Consult your veterinarian for a planned herd health program.
5. When calves are housed in groups after weaning, provide adequate feeding space.