

CO-OP PINNACLE LOW STARCH

For Broodmares, Stallions, Maintenance or Light to Moderate Performance Horses

GUARANTEED ANALYSIS

| | |
|--------------------------------|--------------------|
| Crude Protein (min.) | 12.00% |
| Lysine (min.) | 0.70% |
| Methionine (min.) | 0.25% |
| Crude Fat (min.) | 6.00% |
| Crude Fiber (max.) | 20.00% |
| Acid Detergent Fiber (max.) | 27.00% |
| Neutral Detergent Fiber (max.) | 43.00% |
| Dietary Starch (max.) | 12.00% |
| Sugars (max.) | 3.00% |
| Calcium (min.) | 0.75% (max.) 1.25% |
| Phosphorus (min.) | 0.50% |
| Copper (min.) | 60 ppm |
| Selenium (min.) | 0.50 ppm |
| Zinc (min.) | 150 ppm |
| Vitamin A (min.) | 4,000 IU/lb |
| Vitamin D (min.) | 500 IU/lb |
| Vitamin E (min.) | 125 IU/lb |
| Biotin (min.) | 0.90 mg/lb |

INGREDIENTS

Wheat Middlings, Soybean Hulls, Dehydrated Alfalfa Meal, Flax Meal, Beet Pulp, Rice Mill Feed, Lignin Sulfonate (pelletting agent), Molasses Products, Propionic Acid (a preservative), Calcium Phosphate, Calcium Carbonate, Vegetable Oil with Tertiary Butyl Hydroquinone (a preservative), Salt, dl-Methionine Hydroxy Analog Calcium, Calcium Hydroxide, Zinc Methionine Complex, Manganese Methionine Complex, Copper Lysine Complex, Cobalt Glucoheptonate, Artificial Flavor, Yeast Culture, Vitamin E Supplement, Sodium Selenite, Ascorbic Acid, Selenium Yeast, Biotin, Choline Chloride, d-Calcium Pantothenate, Vitamin A Supplement, Vitamin D-3 Supplement, Thiamine Mononitrate, Niacin Supplement, Riboflavin Supplement, Folic Acid, Vitamin B-12 Supplement, Copper Sulfate, Zinc Sulfate, Zinc Oxide, Manganous Oxide, Manganese Sulfate, Cobalt Carbonate, Calcium Iodate, Acetic Acid, Ammonium Hydroxide, Magnesium Aluminosilicate, Bentonite, Sorbic Acid (a preservative).

FEEDING DIRECTIONS: Co-op Pinnacle Low Starch is designed for horses with a sensitivity to high levels of starch. Follow recommended feeding rates below. Feed similar amounts at the same time each day to minimize digestive upset. Monitor the horse's body condition and adjust feeding practices to maintain an appropriate level of condition. If feeding a grain ration at a rate of 0.50 lb per 100 lb of body weight per day or more, divide total amount into 2 or 3 individual meals offered over 10 to 12 hours. Avoid abrupt changes in feed or forage type. Implement any changes to the diet over a 7- to 10-day transitional period. Changes in the amount of a grain ration should not exceed 1.00 to 1.50 lb per day. Provide clean, fresh water at all times. Consult your Co-op feed representative for specific ration balancing.

| Class | Feeding Rate* |
|--|-----------------|
| Mature, Maintenance | 0.50 to 0.75 lb |
| Mature, Light Work | 0.75 to 1.00 lb |
| Mature, Moderate Work | 1.00 to 1.25 lb |
| Pregnant Mares, first 2/3 of pregnancy | 0.50 to 0.75 lb |
| Non-breeding Stallions | 0.60 to 0.85 lb |
| Breeding Stallions | 0.85 to 1.25 lb |

* Feeding rates are expressed as lb of feed per 100 lb of body weight per day. All calculations assume a daily minimum intake of 1.20 lb of good-quality forage per 100 lb of body weight.

Manufactured By
PROTRITION FEED, LLC
LAVERGNE, TENNESSEE 37086

ITEM NO. 319
TAG CODE NO. 102021-319

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

NET WEIGHT 50 lb (22.68 kg)/INVOICE

319