

# CO-OP PINNACLE SENIOR

For Older Horses with Poor Teeth or Reduced Overall Nutrient Absorption

## GUARANTEED ANALYSIS

Crude Protein (min.)	14.00%
Lysine (min.)	0.65%
Methionine (min.)	0.20%
Crude Fat (min.)	7.00%
Crude Fiber (max.)	18.00%
Acid Detergent Fiber (max.)	21.00%
Neutral Detergent Fiber (max.)	39.00%
Calcium (min.)	0.75%
Phosphorus (min.)	0.60%
Copper (min.)	65 ppm
Selenium (min.)	0.50 ppm
Zinc (min.)	175 ppm
Vitamin A (min.)	5,500 IU/lb
Vitamin D (min.)	500 IU/lb
Vitamin E (min.)	150 IU/lb
Biotin (min.)	0.85 mg/lb

323

## INGREDIENTS

Wheat Middlings, Corn Distillers Dried Grains with Solubles, Soybean Hulls, Rice Mill Feed, Molasses Products, Dehydrated Alfalfa and Timothy, Vegetable Oil, Beet Pulp, Calcium Carbonate, Flax Meal, Salt, Dehydrated Alfalfa Meal, Calcium Phosphate, Dehulled Soybean Meal, Yeast Culture, L-Lysine, Vitamin E Supplement, Artificial Flavor, Sodium Selenite, dl-Methionine Hydroxy Analog Calcium, Calcium Hydroxide, Ascorbic Acid, Zinc Methionine Complex, Manganese Methionine Complex, Copper Lysine Complex, Cobalt Glucoheptonate, Selenium Yeast, Choline Chloride, Biotin, d-Calcium Pantothenate, Vitamin A Supplement, Vitamin D-3 Supplement, Thiamine Mononitrate, Niacin Supplement, Riboflavin Supplement, Folic Acid, Vitamin B-12 Supplement, Copper Sulfate, Zinc Sulfate, Zinc Oxide, Manganous Oxide, Manganese Sulfate, Cobalt Carbonate, Calcium Iodate, Acetic Acid, Ammonium Hydroxide, [Propionic Acid, Sorbic Acid, Tertiary Butyl Hydroquinone (preservatives)].

**FEEDING DIRECTIONS:** Co-op Pinnacle Senior is designed to be a nutritionally dense complete feed for older horses or those that do not consume forage well. As a complete feed, Co-op Pinnacle Senior can replace all or some of the forage in a horse's diet. Follow recommended feeding rates below. If hay is provided in addition to Co-op Pinnacle Senior, limit daily hay intake to 0.5 lb per 100 lb of body weight. Feed similar amounts at the same time each day to minimize digestive upset. Monitor the horse's body condition and adjust feeding practices to maintain an appropriate level of condition. If feeding a grain ration at a rate of 0.50 lb per 100 lb of body weight per day or more, divide total amount into 2 or 3 individual meals offered over 10 to 12 hours. Avoid abrupt changes in feed or forage type. Implement any changes to the diet over a 7 to 10 day transitional period. Changes in the amount of a grain ration should not exceed 1.00 to 1.50 lb per day. Provide clean, fresh water at all times.

Class	Feeding Rate*
Mature, Maintenance	1.25 to 2.00 lb
Mature, Light Work	1.25 to 2.00 lb
Mature, Moderate Work	1.75 to 2.00 lb
Mature, Heavy Work	2.00 to 2.25 lb
Pregnant Mares, first 2/3 of pregnancy	1.25 to 2.00 lb
Pregnant Mares, last 1/3 of pregnancy	1.50 to 2.00 lb
Non-breeding Stallions	1.25 to 2.00 lb
Breeding Stallions	1.25 to 2.00 lb

\* Feeding rates are expressed as lb of feed per 100 lb of body weight per day.

Manufactured By  
PROTRITION FEED, LLC  
LAVERGNE, TENNESSEE 37086

ITEM NO. 323

TAG CODE NO. 080921-323

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

NET WEIGHT 50 lb (22.68 kg)/INVOICE