## CO-OP PINNACLE MARE AND FOAL

For Broodmares, Foals and Growing Horses Up To 2 Years of Age CHAPANTEED ANALYSIS

	D ANALTOIO
Crude Protein (min.)	16.00%
Lysine (min.)	
Methionine (min.)	0.27%
Crude Fat (min.)	6.00%
Methionine (min.) Crude Fat (min.) Crude Fiber (max.)	10.00%
Acid Detergent Fiber (max.)	15.00%
Neutral Detergent Fiber (max.)	
Calcium (min.) 1.00%	(max.) 1.50%
Calcium (min.)	
Copper (min.)	65 ppm
Selenium (mín.)	0.65 ppm
Zinc (min.)	200 ppm
Vitamin A (min.)	10.000 lÚ/lb
Vitamin D (min.) Vitamin E (min.)	1.000 IU/lb
Vitamin E (min.)	
Biotin (min.)	1.00 mg/lb



## INGREDIENTS

Oats, Dehulled Soybean Meal, Molasses Products, Rice Mill Feed, Beet Pulp, Corn Distillers Dried Grains with Solubles, Vegetable Oil, Wheat Middlings, Cracked Corn, Calcium Carbonate, Calcium Phosphate, Flax Meal, Salt, L-Lysine, Cottonseed Meal, Vitamin E Supplement, dl-Methionine, Lignin Sulfonate, Yeast Culture, Potassium Sulfate, Magnesium Sulfate, Ascorbic Acid, Yucca schidigera Extract, Vitamin A Supplement, Vitamin D-3 Supplement, Sodium Selenite, Selenium Yeast, Zinc Methionine Complex, Manganese Methionine Complex, Copper Lysine Complex, Cobalt Glucoheptonate, Choline Chloride, Biotin, d-Calcium Pantothenat, Thiamine Mononitrate, Niacin Supplement, Riboflavin Supplement, Folic Acid, Vitamin B-12 Supplement, Copper Sulfate, Zinc Sulfate, Zinc Oxide, Manganous Oxide, Manganese Sulfate, Cobalt Carbonate, Calcium Iodate, Magnesium Oxide, Ferrous Sulfate, Ethylenediamine Dihydriodide, Acetic Acid, Ammonium Hydroxide, Magnesium Aluminosilicate, Bentonite, [Propionic Acid, Sorbic Acid, Tertiary Butyl Hydroquinone, (preservatives)].

FEEDING DIRECTIONS: Co-op Pinnacle Mare and Foal is designed to provide high-quality protein and energy to broodmares and young growing foals up to 2 years of age. Follow recommended feeding rates below. Feed similar amounts at the same time each day to minimize digestive upset. Monitor the horse's body condition and adjust feeding practices to maintain an appropriate level of condition. If feeding a grain ration at a rate of 0.50 lb per 100 lb of body weight per day or more, divide total amount into 2 or 3 individual meals offered over 10 to 12 hours. Avoid abrupt the processing the changes in feed or forage type. Implement any changes to the diet over a 7- to 10-day transitional period. Changes in the amount of a grain ration should not exceed 1.00 to 1.50 lb per day. Provide clean, fresh water at all times.

•	' '
Class	Feeding Rate*
Barren Mares / Open Mares	0.25 to 0.50 lb
Pregnant Mares, first 2/3 of pregnancy	0.25 to 0.50 lb
Pregnant Mares, last 1/3 of pregnancy	0.50 to 1.00 lb
Lactating Mares, from foaling to 3 months	1.00 to 1.50 lb
Lactating Mares, from 3 months to weaning	0.75 to 1.25 lb
Nursing Foals	Creep Feed
Weaned Foals	0.50 to 1.00 lb
Yearlings & 2-year-olds	0.75 to 1.00 lb

<sup>\*</sup> Feeding rates are expressed as lb of feed per 100 lb of body weight per day. All calculations assume a daily minimum intake of 1.20 lb of good-quality forage per 100 lb of body weight.

Manufactured By PROTRITION FEED, LLC LAVERGNE, TENNESSÉE 37086

**ITEM NO. 331** TAG CODE NO. 070721-331

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.