

# CO-OP PINNACLE BALANCER

For Young Growing Horses, Broodmares, Stallions, Mature Maintenance and Performance Horses

## GUARANTEED ANALYSIS

Crude Protein (min.)	28.00%
Lysine (min.)	2.50%
Methionine (min.)	0.60%
Crude Fat (min.)	3.00%
Crude Fiber (max.)	8.00%
Acid Detergent Fiber (max.)	14.00%
Neutral Detergent Fiber (max.)	25.00%
Calcium (min.)	3.00% (max.)
Phosphorus (min.)	1.50%
Copper (min.)	200 ppm
Selenium (min.)	2 ppm
Zinc (min.)	500 ppm
Vitamin A (min.)	24,000 IU/lb
Vitamin D (min.)	3,000 IU/lb
Vitamin E (min.)	500 IU/lb
Biotin (min.)	4 mg/lb

336

## INGREDIENTS

Dehulled Soybean Meal, Wheat Middlings, Corn Distillers Dried Grains with Solubles, Rice Mill Feed, Calcium Carbonate, Calcium Phosphate, Lignin Sulfonate (pelletizing agent), Dehydrated Alfalfa Meal, L-Lysine, Flax Meal, Cottonseed Meal, Salt, Molasses Products, Yeast Culture, dl-Methionine Hydroxy Analog Calcium, Calcium Hydroxide, Vitamin E Supplement, Sodium Selenite, Artificial Flavor, Zinc Methionine Complex, Manganese Methionine Complex, Copper Lysine Complex, Cobalt Glucoheptonate, Ascorbic Acid, Selenium Yeast, Choline Chloride, Biotin, d-Calcium Pantothenate, Vitamin A Supplement, Vitamin D-3 Supplement, Thiamine Mononitrate, Niacin Supplement, Riboflavin Supplement, Folic Acid, Vitamin B-12 Supplement, Copper Sulfate, Zinc Sulfate, Zinc Oxide, Manganous Oxide, Manganese Sulfate, Cobalt Carbonate, Calcium Iodate, Acetic Acid, Ammonium Hydroxide, Magnesium Aluminosilicate, Bentonite, [Propionic Acid, Sorbic Acid (preservatives)].

**FEEDING DIRECTIONS:** Co-op Pinnacle Balancer is designed to be fed alone to fortify a forage-only diet or mixed with grains to create a balanced concentrate. Follow recommended feeding rates below. Feed similar amounts at the same time each day to minimize digestive upset. Monitor the horse's body condition and adjust feeding practices to maintain an appropriate level of condition. If feeding a grain ration at a rate of 0.50 lb per 100 lb of body weight per day or more, divide total amount into 2 or 3 individual meals offered over 10 to 12 hours. Avoid abrupt changes in feed or forage type. Implement any changes to the diet over a 7- to 10-day transitional period. Changes in the amount of a grain ration should not exceed 1.00 to 1.50 lb per day. Provide clean, fresh water at all times.

### Pinnacle Balancer (FED ALONE):

Class	Feeding Rate*
Mature, Maintenance	0.10 to 0.20
Mature, Light Work	0.15 to 0.25
Non-breeding Stallions	0.10 to 0.25

\* Feeding Rates are expressed as lb of feed per 100 lb of body weight per day. All calculations assume a daily minimum intake of 1.20 lb of good-quality forage per 100 lb of body weight.

### MIXING DIRECTIONS:

For a more energy-dense feed, Pinnacle Balancer can be mixed with oats to create a nutritionally balanced ration. The table to the right provides mixing guidelines for creating a 12% protein, 14% protein, or 16% protein feed. For example, mix 45 lb of oats with 5 lb of Pinnacle Balancer to create 50 lb of a 12% protein feed. The feeding table below provides general recommendation for the 12%, 14%, or 16% feeds.	Protein Level	For a 50 lb Batch of Feed:		
		12% Protein Feed	14% Protein Feed	16% Protein Feed
	Oats	45 lb	40 lb	35 lb
	Pinnacle Balancer	5 lb	10 lb	15 lb
	Final Feed Weight	50 lb	50 lb	50 lb

Class	Feeding Rate*		
	12% Protein Feed	14% Protein Feed	16% Protein Feed
Mature, Maintenance	0.50 to 0.75	0.25 to 0.50	Not Optimal
Mature, Light Work	0.50 to 0.75	0.50 to 0.75	Not Optimal
Mature, Moderate Work	0.75 to 1.25	0.75 to 1.25	0.75 to 1.25
Mature, Heavy Work	1.00 to 1.25	1.00 to 1.50	0.50 to 1.00
Pregnant Mares, first 2/3 of pregnancy	0.25 to 0.75	0.25 to 0.75	0.25 to 0.50
Pregnant Mares, last 1/3 of pregnancy	Not Optimal	0.75 to 1.00	0.50 to 1.00
Lactating Mares (from foaling to 3 months)	Not Optimal	1.25 to 1.50	1.00 to 1.50
Lactating Mares (from 3 months to weaning)	Not Optimal	1.25 to 1.50	0.75 to 1.25
Non-breeding Stallions	0.50 to 0.75	0.40 to 0.60	Not Optimal
Breeding Stallions	0.75 to 1.00	0.60 to 1.00	Not Optimal
Nursing Foals	Not Optimal	Creep Feed	Creep Feed
Weaned Foals	Not Optimal	0.75 to 1.25	0.50 to 1.00
Yearlings & 2-year-olds	Not Optimal	0.75 to 1.25	0.75 to 1.25

\* Feeding rates are expressed as lb of feed per 100 lb of body weight per day. All calculations assume a daily minimum intake of 1.20 lb of good-quality forage per 100 lb of body weight.

ITEM NO. 336

TAG CODE NO. 080921-336

Manufactured By  
**PROTRITION FEED, LLC**  
**LAVERGNE, TENNESSEE 37086**

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

**NET WEIGHT 50 lb (22.68 kg)/INVOICE**