

# CO-OP 48% BEEF CONCENTRATE - 360 RUM MEDICATED

For increased rate of weight gain in growing cattle on pasture or in dry lot (stocker and feeder; dairy and beef replacement heifers).

## ACTIVE DRUG INGREDIENT

Monensin ..... 360 g/ton

## GUARANTEED ANALYSIS

Crude Protein (min.) .....	48.00%
(This includes not more than 24.00% equivalent crude protein from non-protein nitrogen.)	
Crude Fat (min.) .....	2.00%
Crude Fiber (max.) .....	10.00%
Calcium (min.) .....	2.50% (max.) ..... 3.50%
Phosphorus (min.) .....	1.00%
Salt (min.) .....	6.00% (max.) ..... 7.00%
Potassium (min.) .....	1.00%
Sulfur (min.) .....	0.40%
Vitamin A (min.) .....	37,500 IU/lb

# 446

## INGREDIENTS

Plant Protein Products, Processed Grain By-Products, Urea, Salt, Calcium Carbonate, Animal Protein Products, Molasses Products, Magnesium Sulfate, Potassium Sulfate, Sodium Selenite, Calcium Phosphate, Magnesium Oxide, Copper Sulfate, Zinc Oxide, Zinc Sulfate, Manganous Oxide, Manganese Sulfate, Ferrous Sulfate, Cobalt Carbonate, Ethylenediamine Dihydriodide, Calcium Iodate, Vitamin A Acetate, Vitamin D-3 Supplement, Vitamin E Supplement.

**FEEDING DIRECTIONS:** Feed 0.28 to 1.11 lb per head per day thoroughly mixed into not less than 1 lb of the daily ration to provide 50 to 200 mg of Monensin per head per day. During the first 5 days of feeding, cattle should receive 0.28 to 0.55 lb per head per day thoroughly mixed into not less than 1 lb of the daily ration to provide 50 to 100 mg of Monensin per head per day.

**CAUTION:** Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feeds are safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. **USE AS DIRECTED.**

**Manufactured By  
PROTRITION FEED, LLC  
LAVERGNE, TENNESSEE 37086**

ITEM NO. 446

TAG CODE NO. 062310-446

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

**NET WEIGHT 50 lb (22.68 kg)/INVOICE**