

CO-OP 60% BEEF CONCENTRATE - RUM

MEDICATED

For Weaned Cattle on Pasture

For increased rate of weight gain in growing cattle on pasture or in dry lot (stocker and feeder; dairy and beef replacement heifers).

ACTIVE DRUG INGREDIENT

Monensin 304 g/ton

GUARANTEED ANALYSIS

Crude Protein (min.) 60.00%
 (This includes not more than 30.00% equivalent crude protein from non-protein nitrogen.)
 Crude Fat (min.) 1.00%
 Crude Fiber (max.) 10.00%
 Calcium (min.) 5.00% (max.) 6.00%
 Phosphorus (min.) 1.00%
 Salt (min.) 7.00% (max.) 8.00%
 Potassium (min.) 1.00%
 Zinc (min.) 900 ppm
 Vitamin A (min.) 50,000 IU/lb
 Vitamin E (min.) 140 IU/lb

482

INGREDIENTS

Plant Protein Products, Animal Protein Products, Urea, Calcium Carbonate, Salt, Magnesium Sulfate, Potassium Sulfate, Molasses Products, Sodium Selenite, Lignin Sulfonate, Zinc Oxide, Zinc Sulfate, Manganese Sulfate, Manganous Oxide, Ferrous Sulfate, Magnesium Oxide, Copper Sulfate, Cobalt Carbonate, Ethylenediamine Dihydriodide, Calcium Iodate, Vitamin A Acetate, Vitamin D-3 Supplement, Vitamin E Supplement.

FEEDING DIRECTIONS: Co-op 60% Beef Concentrate-RUM is to be fed at a rate to provide 50 to 200 mg of Monensin per head per day. During the first 5 days of feeding, cattle should receive 50 to 100 mg of Monensin per head per day. The daily allowance of Monensin must be provided in not less than 1 lb of feed. Use the mixing chart below to blend Co-op 60% Beef Concentrate-RUM with corn. Feed at the rates indicated in the left column to provide 50 to 200 mg of Monensin per head per day. Be sure adequate pasture is available.

LB OF FEED HEAD/DAY	LB OF 60% BEEF CONCENTRATE-RUM	LB OF CORN	PROTEIN% OF MIX
6.6 - 26.3	100	1900	11%
4.4 - 17.5	150	1850	12%
3.0 - 11.6	225	1775	14%
2.2 - 8.7	300	1700	16%

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feeds are safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin could be fatal to cattle. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. USE AS DIRECTED.

Manufactured By
PROTRITION FEED, LLC
 LAVERGNE, TENNESSEE 37086

ITEM NO. 482
 TAG CODE NO. 062310-482

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

NET WEIGHT 50 lb (22.68 kg)/INVOICE