

CO-OP PUNCH 13 25-MEC

MEDICATED

Starter for Weaned Pigs

For control of swine dysentery (vibronic dysentery, bloody scours or hemorrhagic dysentery); control of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by *Salmonella choleraesuis*); for increased rate of weight gain and improvement of feed efficiency.

ACTIVE DRUG INGREDIENT

Carbadox 50 g/ton

GUARANTEED ANALYSIS

Crude Protein (min.)	20.00%
Lysine (min.)	1.30%
Crude Fat (min.)	5.00%
Crude Fiber (max.)	4.00%
Calcium (min.) 0.75% (max.)	1.25%
Phosphorus (min.)	0.70%
Salt (min.) 0.30% (max.)	0.80%
Selenium (min.)	0.30 ppm
Zinc (min.)	2,500 ppm

502

INGREDIENTS

Ground Corn, Dehulled Soybean Meal, Dried Whey, Fish Meal, Animal Fat and Vegetable Oil, Sugar, Calcium Phosphate, Calcium Carbonate, Zinc Oxide, Copper Sulfate, L-Lysine, Natural and Artificial Flavors, Yucca Schidegera Extract, Choline Chloride, Vitamin E Supplement, Niacin Supplement, Calcium Pantothenate, Biotin, Menadione Sodium Bisulfite Complex, Riboflavin Supplement, Vitamin A Acetate, Vitamin B-12 Supplement, Vitamin D-3 Supplement, Folic Acid, Manganous Oxide, Ferrous Sulfate, Calcium Iodate, Cobalt Carbonate, Sodium Selenite, Ethoxyquin (a preservative).

FEEDING DIRECTIONS: Feed as sole ration until two weeks after weaning or until pigs weigh 25 lb whichever occurs first. Always provide plenty of clean, fresh water in clean waterers.

WARNING: Do not feed to swine within 42 days before slaughter.

CAUTION: Not for use in pregnant swine or swine intended for breeding purposes.

Manufactured By
PROTRITION FEED, LLC
LAVERGNE, TENNESSEE 37086

ITEM NO. 502

TAG CODE NO. 110316-502

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

NET WEIGHT 50 lb (22.68 kg)/INVOICE