

CO-OP VICTORY LANE CALF STARTER-DEC CL MEDICATED

For Calves Weighing Up To 250 lb

For the prevention of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii*.

ACTIVE DRUG INGREDIENT

Decoquinatate 45.4 g/ton

For the prevention of adult house and stable fly emergence from the manure of treated cattle.

ACTIVE INGREDIENT:

Diflubenzuron (CAS #35367-38-5) 0.001000%

OTHER INGREDIENTS 99.999000%

Total 100.000000%

GUARANTEED ANALYSIS

Crude Protein (min.) 20.50%

Crude Fat (min.) 3.50%

Crude Fiber (max.) 8.00%

Acid Detergent Fiber (max.) 11.80%

Calcium (min.) 1.25% (max.) 1.75%

Phosphorus (min.) 0.77%

Salt (min.) 0.70% (max.) 1.20%

Potassium (min.) 0.80%

Selenium (min.) 0.45 ppm

Vitamin A (min.) 10,000 IU/lb

INGREDIENTS

Grain Products, Plant Protein Products, Processed Grain By-Products, Molasses Products, Calcium Carbonate, Calcium Phosphate, Propionic Acid (a preservative), Animal Protein Products, Vegetable Oil with Tertiary Butyl Hydroquinone (a preservative), Salt, dl-Methionine Hydroxy Analog Calcium, Calcium Hydroxide, L-Lysine, Selenium Yeast, Sodium Selenite, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Magnesium Oxide, Copper Sulfate, Zinc Oxide, Zinc Sulfate, Manganous Oxide, Manganese Sulfate, Ferrous Sulfate, Cobalt Carbonate, Ethylenediamine Dihydroiodide, Calcium Iodate, Acetic Acid, Sodium Propionate, Ammonium Hydroxide, Sorbic Acid (a preservative), Magnesium Aluminosilicate.

FEEDING DIRECTIONS: Feed Co-op Victory Lane Calf Starter-DEC CL by 2 days of age at the rate of 1.0% of body weight per day to provide 22.7 mg of Decoquinatate per 100 lb of body weight per day and 0.1 mg of Diflubenzuron per kg of body weight per day. Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates that coccidiosis is likely to be a hazard. Do not exceed 0.30 ppm supplemental selenium in the total ration. Always provide plenty of clean, fresh water.

WARNING: Do not feed to cows producing milk for human consumption.

CAUTION: Do not use Decoquinatate in feeds containing bentonite. Do not feed to sheep or other copper-sensitive animals.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

Causes moderate eye injury. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield or safety glasses). Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse. Wear long-sleeved shirt and long pants; socks, shoes and gloves.

Storage and Disposal: Do not contaminate water, food or feed by storage or disposal.

Storage: Store pesticide in original container in a dry place.

Product Disposal: Wastes resulting from the use of this product may be disposed of onsite or at an approved waste disposal facility.

Container Disposal: Completely empty bag. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

When used as directed, this product will provide sufficient Diflubenzuron larvacide to prevent the emergence of adult house and stable flies from manure of treated cattle. Existing adult flies will not be affected. Start feeding before house and stable flies appear and continue use until cold weather marks the end of the fly season. Feed this product at the rate of 1 lb per 100 lb of body weight per day which will provide 0.1 mg of Diflubenzuron per kg of body weight per day. Introduction of this product after an adult house or stable fly infestation is established will require treatment of cattle and confinement areas with adulticides if immediate elimination of the adult fly population is desired.

Manufactured By
PROTRITION FEED, LLC
LAVERGNE, TENNESSEE 37086

ITEM NO. 93085

TAG CODE NO. 022122-93085

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

NET WEIGHT 50 lb (22.68 kg)/INVOICE

93085