CO-OP CALF PRIMER/TCR I-MON CL

MEDICATED

Starter for Growing Calves
For the prevention and control of coccidiosis caused by Eimeria bovis and Eimeria zuernii.

ACTIVE DRUG INGREDIENT

For the prevention of adult house and stable fly emergence from the manure of treated cattle. ACTIVE INGREDIENT: | Diflubenzuron (CAS #35367-38-5) | 0.000333% | OTHER INGREDIENTS | 99.999667%100.000000% Total **GUARANTEED ANALYSIS**
 Crude Fat (min.)
 2.00%

 Crude Fiber (max.)
 15.00%

 Acid Detergent Fiber (max.)
 21.00%

 Calcium (min.)
 0.70%
 (max.)
 1.20%



WARNING: A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

CAUTION: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feeds are safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. Do not feed to sheep or other copper-sensitive animals.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS
Causes moderate eye injury. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse. Wear long-sleeved shirt and long pants, socks, shoes, and gloves.

Storage and Disposal: Do not contaminate water, food or feed by storage or disposal.

Storage: Store pesticide in original container in a dry place.

Product Disposal: Wastes resulting from the use of this product may be disposed of onsite or at an approved waste disposal facility.

Container Disposal: Completely empty bag. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

When used as directed, this product will provide sufficient Difflubenzuron larvicide to prevent the emergence of adult house and stable flies from manure of treated cattle. Existing adult flies will not be affected. Start feeding before house and stable flies appear and continue use until cold weather marks the end of the fly season. Feed this product at the rate of 3 lb per 100 lb of body weight per day which will provide 0.1 mg of Diffubenzuron per kg of body weight per day. Introduction of this product after an adult house or stable fly infestation is established will require treatment of cattle and confinement areas with adulticides if immediate elimination of the adult fly population is desired.

Manufactured By PROTRITION FEED, LLC **LAVERGNE, TENNESSEE 37086**

ITEM NO. 93115 TAG CODE NO. 030123-93115

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.