CO-OP 17% PELLETED SHOW GOAT GROWER-MON

MEDICATED

For Growing and Finishing Goats Being Fitted for Show For the prevention of coccidiosis caused by *Eimeria crandallis*, *Eimeria christenseni* and *Eimeria ninakohlyakimovae*.

	ACTIVE DR	RUG INGREDIE	NT	
Monensin			20 g/ton	
	GUARANT	EED ANALYSI	s	
Crude Protein (min.)			17.00%	$\overline{}$
(This includes not more than 1.2				•
Crude Fat (min.)			4.00%	\
Crude Fiber (max.)				
Acid Detergent Fiber (max.)			20.00%	M
Calcium (min.)	0.75%	(max.)	1.25%	_
Phosphorus (min.)			0.40%	
Salt (min.)				М
Potassium (min.)			1.10%	
Copper (min.)	20 ppm	(max.)	27 ppm	
Selenium (min.)				
Zinc (min.)				
Vitamin A (min.)				
Vitamin E (min.)				
. ,				

INGREDIENTS

Wheat Middlings, Ground Corn, Dehydrated Alfalfa Meal, Corn Distillers Dried Grains with Solubles, Dehulled Soybean Meal, Soybean Hulls, Cottonseed Hulls, Dried Whey, Molasses Products, Propionic Acid (a preservative), Calcium Carbonate, Lignin Sulfonate, Salt, Vegetable Oil with Tertiary Butyl Hydroquinone (a preservative), Ammonium Chloride, Animal Fat and Vegetable Oil with Ethoxyquin (a preservative), Yeast Culture, Selenium Yeast, Zinc Methionine Complex, Manganese Methionine Complex, Copper Lysine Complex, Cobalt Glucoheptonate, Vitamin A Supplement, Vitamin D-3 Supplement, Natural and Artificial Flavors, Dehydrated Lactobacillus acidophilus Fermentation Product, Dehydrated Lactobacillus casei Fermentation Product, Dehydrated Bifidobacterium thermophilum Fermentation Product, Dehydrated Enterococcus faecium Fermentation Product, Vitamin E Supplement, Zinc Sulfate, Sodium Molybdate, Zinc Oxide, Ferrous Sulfate, Manganese Sulfate, Sodium Selenite, Manganous Oxide, Ethylenediamine Dihydriodide, Cobalt Carbonate.

FEEDING DIRECTIONS: Feed continuously as the sole ration to growing goats 6 weeks of age or older. Do not allow animals to run out of feed.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin-medicated goat feed is safe for use in goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result. Do not feed to lactating goats. Do not feed to sheep or other copper-sensitive animals.

Manufactured By PROTRITION FEED, LLC LAVERGNE, TENNESSEE 37086

ITEM NO. 93326

TAG CODE NO. 103122-93326

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

NET WEIGHT 50 lb (22.68 kg)/INVOICE