

# CO-OP PINNACLE LOW STARCH

For Broodmares, Stallions, Maintenance or Light to Moderate Performance Horses

## GUARANTEED ANALYSIS

Crude Protein (min.) .....	12.00%
Lysine (min.) .....	0.70%
Methionine (min.) .....	0.25%
Crude Fat (min.) .....	6.00%
Crude Fiber (max.) .....	20.00%
Acid Detergent Fiber (max.) .....	27.00%
Neutral Detergent Fiber (max.) .....	43.00%
Dietary Starch (max.) .....	10.00%
Calcium (min.) .....0.75% (max.).....	1.25%
Phosphorus (min.) .....	0.50%
Copper (min.) .....	60 ppm
Selenium (min.) .....	0.50 ppm
Zinc (min.) .....	150 ppm
Vitamin A (min.) .....	4,000 IU/lb
Vitamin D (min.) .....	500 IU/lb
Vitamin E (min.) .....	125 IU/lb
Biotin (min.) .....	0.90 mg/lb

## INGREDIENTS

Soybean Hulls, Wheat Middlings, Dehydrated Alfalfa Meal, Flax Meal, Beet Pulp, Rice Bran, Lignin Sulfonate (pelleting agent), Molasses Products, Calcium Carbonate, Salt, Calcium Phosphate, Vegetable Oil, dl-Methionine Hydroxy Analog Calcium, Calcium Hydroxide, Zinc Methionine Complex, Manganese Methionine Complex, Copper Lysine Complex, Cobalt Glucoheptonate, Artificial Flavor, Yeast Culture, Vitamin E Supplement, Sodium Selenite, Ascorbic Acid, Biotin, Choline Chloride, Calcium Pantothenate, Vitamin A Supplement, Vitamin D-3 Supplement, Thiamine Mononitrate, Niacin Supplement, Riboflavin Supplement, Folic Acid, Vitamin B-12 Supplement, Copper Sulfate, Ferrous Sulfate, Manganese Oxide, Zinc Sulfate, Zinc Oxide, Selenium Yeast, Cobalt Carbonate, Calcium Iodate, Acetic Acid, Ammonium Hydroxide, Magnesium Aluminosilicate, Bentonite, [Propionic Acid, Sorbic Acid, Tertiary Butyl Hydroquinone (preservatives)].

**FEEDING DIRECTIONS:** Co-op Pinnacle Low Starch is designed for horses with a sensitivity to high levels of starch. Follow recommended feeding rates below. Feed similar amounts at the same time each day to minimize digestive upset. Monitor the horse's body condition and adjust feeding practices to maintain an appropriate level of condition. If feeding a grain ration at a rate of 0.50 lb per 100 lb of body weight per day or more, divide total amount into 2 or 3 individual meals offered over 10 to 12 hours. Avoid abrupt changes in feed or forage type. Implement any changes to the diet over a 7 to 10 day transitional period. Changes in the amount of a grain ration should not exceed 1.00 to 1.50 lb per day. Provide clean, fresh water at all times.

Class	Feeding Rate*
Mature, Maintenance	0.50 to 0.75 lb
Mature, Light Work	0.75 to 1.00 lb
Mature, Moderate Work	1.00 to 1.25 lb
Pregnant Mares, first 2/3 of pregnancy	0.50 to 0.75 lb
Non-breeding Stallions	0.60 to 0.85 lb
Breeding Stallions	0.85 to 1.25 lb

\* Feeding rates are expressed as lb of feed per 100 lb of body weight per day. All calculations assume a daily minimum intake of 1.20 lb of good-quality forage per 100 lb of body weight.

Manufactured By  
**TENNESSEE FARMERS COOPERATIVE**  
 LAVERGNE, TENNESSEE 37086

ITEM NO. 319  
 TAG CODE NO. 111419-319

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

**NET WEIGHT 50 lb (22.68 kg)/INVOICE**

- JU  
 - LOT

# 319

