

CO-OP PINNACLE 1000

For Broodmares, Stallions, Mature Maintenance and Performance Horses

GUARANTEED ANALYSIS

Crude Protein (min.)	10.00%	
Lysine (min.)	0.60%	
Methionine (min.)	0.20%	
Crude Fat (min.)	5.00%	
Crude Fiber (max.)	10.00%	
Acid Detergent Fiber (max.).....	15.00%	
Neutral Detergent Fiber (max.)	24.00%	
Calcium (min.)	0.50% (max.).....	1.00%
Phosphorus (min.)	0.45%	
Copper (min.)	55 ppm	
Selenium (min.)	0.30 ppm	
Zinc (min.)	140 ppm	
Vitamin A (min.)	5,000 IU/lb	
Vitamin D (min.)	500 IU/lb	
Vitamin E (min.)	100 IU/lb	
Biotin (min.)	0.65 mg/lb	

INGREDIENTS

Oats, Cracked Corn, Molasses Products, Wheat Middlings, Beet Pulp, Rice Bran, Roasted Soybeans, Corn Distillers Dried Grains with Solubles, Rice Hulls, Calcium Carbonate, Calcium Phosphate, Vegetable Oil, Flax Meal, L-Lysine, Salt, Soybean Hulls, Dehulled Soybean Meal, dl-Methionine, Vitamin E Supplement, Sodium Selenite, Cottonseed Meal, Yeast Culture, Lignin Sulfonate, Zinc Sulfate, Ascorbic Acid, Yucca Schidigera Extract, Vitamin A Supplement, Vitamin D-3 Supplement, Zinc Methionine Complex, Manganese Methionine Complex, Copper Lysine Complex, Cobalt Glucoheptonate, Choline Chloride, Biotin, Calcium Pantothenate, Thiamine Mononitrate, Niacin Supplement, Riboflavin Supplement, Folic Acid, Vitamin B-12 Supplement, Copper Sulfate, Ferrous Sulfate, Manganous Oxide, Zinc Oxide, Selenium Yeast, Cobalt Carbonate, Calcium Iodate, Magnesium Oxide, Manganese Sulfate, Ethylenediamine Dihydriodide, Acetic Acid, Ammonium Hydroxide, Magnesium Aluminosilicate, Bentonite, [Propionic Acid, Sorbic Acid, Tertiary Butyl Hydroquinone, (preservatives)].

FEEDING DIRECTIONS: Co-op Pinnacle 1000 is designed to provide high-quality protein and energy to mature horses. Follow recommended feeding rates below. Feed similar amounts at the same time each day to minimize digestive upset. Monitor the horse's body condition and adjust feeding practices to maintain an appropriate level of condition. If feeding a grain ration at a rate of 0.50 lb per 100 lb of body weight per day or more, divide total amount into 2 or 3 individual meals offered over 10 to 12 hours. Avoid abrupt changes in feed or forage type. Implement any changes to the diet over a 7 to 10 day transitional period. Changes in the amount of a grain ration should not exceed 1.0 to 1.5 lb per day. Provide clean, fresh water at all times.

Class	Feeding Rate*
Mature, Maintenance	0.30 to 0.50 lb
Mature, Light Work	0.50 to 0.75 lb
Mature, Moderate Work	0.75 to 1.00 lb
Mature, Heavy Work	1.00 to 1.25 lb
Pregnant Mares, first 2/3 of pregnancy	0.50 to 0.75 lb
Non-breeding Stallions	0.50 to 0.75 lb
Breeding Stallions	0.75 to 1.00 lb

* Feeding rates are expressed as lb of feed per 100 lb of body weight per day. All calculations assume a daily minimum intake of 1.20 lb of good-quality forage per 100 lb of body weight.

Manufactured By
TENNESSEE FARMERS COOPERATIVE
 LAVERGNE, TENNESSEE 37086

ITEM NO. 328
 TAG CODE NO. 111419-328

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

NET WEIGHT 50 lb (22.68 kg)/INVOICE

- JU
 - LOT

328

