


CO-OP PINNACLE 1200

For Young Growing Horses, Broodmares, Stallions, Mature Maintenance and Performance Horses

GUARANTEED ANALYSIS

Crude Protein (min.)	12.00%
Lysine (min.)	0.75%
Methionine (min.)	0.20%
Crude Fat (min.)	6.00%
Crude Fiber (max.)	10.00%
Acid Detergent Fiber (max.)	16.00%
Neutral Detergent Fiber (max.)	27.00%
Calcium (min.)	0.75% (max.) 1.25%
Phosphorus (min.)	0.60%
Copper (min.)	55 ppm
Selenium (min.)	0.45 ppm
Zinc (min.)	200 ppm
Vitamin A (min.)	7,500 IU/lb
Vitamin D (min.)	500 IU/lb
Vitamin E (min.)	150 IU/lb
Biotin (min.)	1.00 mg/lb



INGREDIENTS

Oats, Wheat Middlings, Roasted Soybeans, Molasses Products, Beet Pulp, Corn Distillers Dried Grains with Solubles, Vegetable Oil, Rice Hulls, Cracked Corn, Calcium Carbonate, Calcium Phosphate, Corn Germ Meal, Rice Bran, Flax Meal, L-Lysine, Salt, Soybean Hulls, Dehulled Soybean Meal, dl-Methionine, Sodium Selenite, Lignin Sulfonate, Yeast Culture, *Saccharomyces cerevisiae*, Zinc Sulfate, Ascorbic Acid, Cottonseed Meal, Yucca Schidigera Extract, Zinc Methionine Complex, Manganese Methionine Complex, Copper Lysine Complex, Cobalt Glucoheptonate, Vitamin E Supplement, Biotin, Choline Chloride, Vitamin A Supplement, Calcium Pantothenate, Thiamine Mononitrate, Vitamin D-3 Supplement, Niacin Supplement, Riboflavin Supplement, Vitamin B-12 Supplement, Folic Acid, Copper Sulfate, Ferrous Sulfate, Manganous Oxide, Zinc Oxide, Selenium Yeast, Cobalt Carbonate, Calcium Iodate, Magnesium Oxide, Manganese Sulfate, Ethylenediamine Dihydrochloride, Acetic Acid, Ammonium Hydroxide, Magnesium Aluminosilicate, Bentonite, [Propionic Acid, Sorbic Acid, Tertiary Butyl Hydroquinone, (preservatives)].

FEEDING DIRECTIONS: Co-op Pinnacle 1200 is designed to provide a high-quality balanced ration to young growing, mature maintenance and performance horses. Follow recommended feeding rates below. Feed similar amounts at the same time each day to minimize digestive upset. Monitor the horse's body condition and adjust feeding practices to maintain an appropriate level of condition. If feeding a grain ration at a rate of 0.50 lb per 100 lb of body weight per day or more, divide total amount into 2 or 3 individual meals offered over 10 to 12 hours. Avoid abrupt changes in feed or forage type. Implement any changes to the diet over a 7 to 10 day transitional period. Changes in the amount of a grain ration should not exceed 1.00 to 1.50 lb per day. Provide clean, fresh water at all times.

Class	Feeding Rate*
Mature, Maintenance	0.50 to 0.75 lb
Mature, Light Work	0.50 to 0.75 lb
Mature, Moderate Work	0.75 to 1.00 lb
Mature, Heavy Work	1.00 to 1.25 lb
Pregnant Mares, first 2/3 of pregnancy	0.50 to 0.75 lb
Yearlings & 2 year olds	0.50 to 0.75 lb
Non-breeding Stallions	0.50 to 0.75 lb
Breeding Stallions	0.75 to 1.00 lb

* Feeding rates are expressed as lb of feed per 100 lb of body weight per day. All calculations assume a daily minimum intake of 1.20 lb of good-quality forage per 100 lb of body weight.

Manufactured By
TENNESSEE FARMERS COOPERATIVE
 LAVERGNE, TENNESSEE 37086

ITEM NO. 333
 TAG CODE NO. 031119-333



Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

NET WEIGHT 50 lb (22.68 kg)/INVOICE

~JU
 ~LOT