

# CO-OP 36% TEND-R-LEEN BEEF CONCENTRATE PELLET-RUM

## MEDICATED

For Confined Feedlot Beef Cattle

For improved feed efficiency.

### ACTIVE DRUG INGREDIENTS

Monensin ..... 240 g/ton

### GUARANTEED ANALYSIS

Crude Protein (min.) ..... 36.00%  
 (This includes not more than 12.00% equivalent crude protein from  
 non-protein nitrogen.)  
 Crude Fat (min.) ..... 1.75%  
 Crude Fiber (max.) ..... 10.00%  
 Calcium (min.) ..... 3.00% (max.) ..... 4.00%  
 Phosphorus (min.) ..... 1.00%  
 Salt (min.) ..... 2.50% (max.) ..... 3.50%  
 Potassium (min.) ..... 0.90%  
 Vitamin A (min.) ..... 25,000 IU/lb  
 Vitamin E (min.) ..... 115 IU/lb

### INGREDIENTS

Plant Protein Products, Processed Grain By-Products, Calcium Carbonate, Urea, Salt, Molasses Products, Propionic Acid (a preservative), Calcium Phosphate, Lignin Sulfonate, Sodium Selenite, Animal Fat and Vegetable Oil with Ethoxyquin (a preservative), Vitamin E Supplement, Magnesium Sulfate, Potassium Sulfate, Zinc Oxide, Calcium Iodate, Cobalt Carbonate, Manganous Oxide, Ferrous Oxide, Vitamin A Acetate (in gelatin), D-Activated Animal Sterol (source of Vitamin D-3), a-Tocopherol Acetate (source of Vitamin E), Choline Chloride, Vitamin B-12 Supplement, Riboflavin Supplement, d-Calcium Pantothenate, Niacin Supplement, Thiamine Mononitrate, Pyridoxine Hydrochloride, Biotin, Folic Acid, Mineral Oil, Magnesium Oxide, Copper Sulfate, Zinc Sulfate, Manganese Sulfate, Ferrous Sulfate, Ethylenediamine Dihydrochloride.

**FEEDING DIRECTIONS:** Feed continuously at the rate of 1.5 lb per head per day thoroughly mixed into the amount of shelled corn to be consumed daily to provide 180 mg of Monensin per head per day. Use this ration after cattle have completed the Tend-R-Leen Starter Program. Always provide plenty of clean, fresh water.

**WARNING:** A withdrawal time has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

**CAUTION:** Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feeds are safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. USE AS DIRECTED. Do not feed to sheep or other copper-sensitive animals.

### Manufactured By


**TENNESSEE FARMERS COOPERATIVE  
LAVERGNE, TENNESSEE 37086**

ITEM NO. 441

TAG CODE NO. 110316-441

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

# 441



**NET WEIGHT 50 lb (22.68 kg)/INVOICE**

-JU  
-LOT