

CO-OP 18% PELLETTED BEEF SUPPLEMENT- RUM

MEDICATED

For Weaned Cattle On Pasture

For increased rate of weight gain in growing cattle on pasture or in dry lot (stocker and feeder; dairy and beef replacement heifers). For improved feed efficiency in cattle (mature reproducing beef cows on pasture or in dry lots).

ACTIVE DRUG INGREDIENT

Monensin 63 g/ton

GUARANTEED ANALYSIS

Crude Protein (min.)	18.00%
Crude Fat (min.)	3.00%
Crude Fiber (max.)	8.00%
Calcium (min.) 0.40% (max.)	0.90%
Phosphorus (min.)	0.65%
Salt (min.) 0.80% (max.)	1.30%
Potassium (min.)	0.80%
Selenium (min.)	0.60 ppm
Vitamin A (min.)	5,000 IU/lb
Vitamin E (min.)	17 IU/lb

INGREDIENTS

Processed Grain By-Products, Plant Protein Products, Molasses Products, Propionic Acid, Roughage Products, Calcium Carbonate, Salt, Grain Products, Sodium Selenite, Selenium Yeast, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper Amino Acid Complex, Cobalt Glucoheptonate, Zinc Oxide, Zinc Sulfate, Manganous Oxide, Manganese Sulfate, Ferrous Sulfate, Magnesium Oxide, Copper Sulfate, Cobalt Carbonate, Ethylenediamine Dihydriodide, Calcium Iodate, Vitamin A Acetate, Vitamin D-3 Supplement, Vitamin E Supplement.

FEEDING DIRECTIONS: During the first 5 days of feeding, cattle should receive 1.59 to 3.17 lb per head per day of Co-op 18% Pelletted Beef Supplement-RUM. Following this adjustment period, feed at the rate of 1.59 to 6.35 lb per head per day to supply not less than 50 nor more than 200 mg of Monensin per head per day while on pasture or in dry lots. Do not self feed. Always supply plentiful good quality forage and plenty of fresh water.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feed is safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. Do not feed to sheep or other copper sensitive animals.

Manufactured By
TENNESSEE FARMERS COOPERATIVE
LAVERGNE, TENNESSEE 37086

ITEM NO. 456
TAG CODE NO. 061510-456

456PE



Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

NET WEIGHT 50 lb (22.68 kg)/INVOICE

~JU
-LOT