



**CO-OP
Feeds**

CO-OP PASTURE BLOCK WITH RABON®

For Weaned Cattle on Pasture

Control of fecal flies in manure of treated cattle. Prevents development of face flies, horn flies, house flies and stable flies in the manure of treated cattle.

ACTIVE INGREDIENT: Tetrachlorvinphos: 2-chloro-1-(2,4,5-trichlorophenyl) vinyl dimethyl phosphate 0.31%*
INERT INGREDIENTS: 99.69%
Total 100.00%

*Each pound contains 1.4 g Rabon.

**Refers only to ingredients which are not Larvicidal.

Rabon is a registered trademark of Fermenta Animal Health Co.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Personal Protective Equipment (PPE):

Mixers and Handlers must wear:

- long-sleeved shirt and pants
- shoes and socks
- chemical resistant gloves

User Safety Requirements: Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

CAUTION

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.

FIRST AID

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger, or if available by administering syrup of ipecac. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF ON SKIN: Wash with plenty of water. Get medical attention.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not contaminate water when disposing of equipment wash water.

GUARANTEED ANALYSIS

Crude Protein (min.)	5.50%	
Crude Fat (min.)	0.50%	
Crude Fiber (max.)	10.00%	
Acid Detergent Fiber (max.)	8.00%	
Calcium (min.)	4.50% (max.)	5.40%
Phosphorus (min.)	2.50%	
Salt (min.)	19.00% (max.)	22.50%
Potassium (min.)	0.60%	
Selenium (min.)	2.14 ppm	
Vitamin A (min.)	40,000 IU/lb	
Vitamin D-3 (min.)	5,000 IU/lb	

INGREDIENTS

Processed Grain By-Products, Salt, Molasses Products, Dicalcium Phosphate, Roughage Products, Calcium Carbonate, Plant Protein Products, Lignin Sulfonate, Sodium Selenite, Iron Oxide, Animal Fat and Vegetable Oil with Ethoxyquin (a preservative), Vitamin A Acetate, Vitamin D-3 Supplement, Copper Sulfate, Vitamin E Supplement, Zinc Oxide, Zinc Sulfate, Manganous Oxide, Manganese Sulfate, Ferrous Sulfate, Magnesium Oxide, Cobalt Carbonate, Ethylenediamine Dihydriodide, Calcium Iodate, Mineral Oil.

625

NOTE: Directions for Use, Storage and Disposal, General Precautions and Restrictions, General Information and Application Instructions on back of label.

EPA Reg. No. 9078-12
EPA Est. No. 9078-TN-4 9078-TN-5

ITEM NO. 625
TAG CODE NO. 012803-625

**Manufactured By
TENNESSEE FARMERS COOPERATIVE
200 WALDRON ROAD
LAVERGNE, TENNESSEE 37086**

NET WEIGHT 33 1/3 lb (15.11 kg)

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with this labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in a dry place in original container.

Disposal: Completely empty container into application (mixer/feeder) equipment. Then dispose of empty bag in sanitary landfill or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

GENERAL PRECAUTIONS AND RESTRICTIONS

Do not apply this product in a way that will contact workers or other persons, either directly or through drift.

Only protected handlers may be in the area during application.

GENERAL INFORMATION

Co-op Pasture Block with Rabon® prevents the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle but is not effective against existing adult flies. When starting a feeding program during the fly season, it is desirable to use other control measures to reduce the population to existing adult flies.

In some cases, supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises.

In order to achieve optimum fly control, Co-op Pasture Block with Rabon® should be used in conjunction with other good management and sanitation practices.

Control of fecal flies in manure of treated cattle. Prevents development of face flies, horn flies, house flies and stable flies in the manure of treated cattle.

APPLICATION INSTRUCTIONS

Feed Co-op Pasture Block with Rabon® at the rate of 1/2 lb per 1000 lb of body weight.

Place one block per five head of cattle.

Provide blocks where animals congregate.

Feed blocks continuously. Never allow animals to run out. Add an additional block when each block has been one-half consumed.

Animals should not be starved for minerals before initiation of Co-op Pasture Block with Rabon®, but omit all salt from other feed after program is started. Do not feed free choice sources of salt or minerals containing salt in any form.

Start feeding Co-op Pasture Block with Rabon® early in the spring before flies begin to appear and continue feeding throughout the summer into the fall until cold weather restricts fly activity.

Co-op Pasture Block with Rabon® may be fed up to slaughter and to lactating dairy cows.

NET WEIGHT 33 1/3 lb (15.11 kg)