

CO-OP FOUNDATION CATTLE MINERAL - RUM

MEDICATED

For Weaned Cattle on Pasture

For increased rate of weight gain in growing cattle on pasture or in dry lot (stocker and feeder; dairy and beef replacement heifers).

ACTIVE DRUG INGREDIENT

Monensin 1,200 g/ton

GUARANTEED ANALYSIS

Calcium (min.)	15.00%	(max.)	18.00%
Phosphorus (min.)			3.00%
Salt (min.)	22.00%	(max.)	26.00%
Magnesium (min.)			1.00%
Copper (min.)			1,300 ppm
Selenium (min.)			30 ppm
Zinc (min.)			2,600 ppm
Vitamin A (min.)			225,000 IU/lb
Vitamin D-3 (min.)			17,000 IU/lb
Vitamin E (min.)			220 IU/lb

INGREDIENTS

Calcium Carbonate, Salt, Calcium Phosphate, Molasses Products, Sodium Selenite, Magnesium Oxide, Animal Fat and Vegetable Oil with Ethoxyquin (a preservative), Copper Sulfate, Iron Oxide, Vitamin A Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Selenium Yeast, Artificial Flavor, Ethylenediamine Dihydriodide, Zinc Oxide, Zinc Sulfate, Manganous Oxide, Manganese Sulfate, Ferrous Sulfate, Cobalt Carbonate, Calcium Iodate.

FEEDING DIRECTIONS: Feed at the rate of 100 g (3.53 oz) per head per day in not less than 1 lb of feed to provide 132.38 mg of Monensin per head per day. During the first five days of feeding, cattle should receive no more than 100 mg of Monensin (2.66 oz of Co-op Foundation Cattle Mineral-RUM) per head per day contained in not less than 1 lb of feed. Feed this product along with ample forage and/or other appropriate feedstuffs. Provide clean, fresh water at all times.

WARNING: A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

CAUTION: Do not allow horses or other equines access to feed containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin-medicated cattle feeds are safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Do not exceed the levels of Monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. Do not feed to sheep or other copper-sensitive animals.

Manufactured By
TENNESSEE FARMERS COOPERATIVE
LAVERGNE, TENNESSEE 37086

ITEM NO. 651

TAG CODE NO. 081817-651

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

NET WEIGHT 50 lb (22.68 kg)/INVOICE

-JU
 -LOT

651

