CO-OP SHOW PIG PRIMER-DEN

MEDICATED

For Growing Show Pigs From 50 to 180 lb

Control of swine dysentery associated with *Brachyspira* (formerly *Serpulina* or *Treponema*) *hyodysenteriae* susceptible to Tiamulin. Control of porcine proliferative enteropathies (ileitis) associated with *Lawsonia intracellularis*.

95002

ACTIVE DRUG INGREDIENT

Tiamulin Hydrogen Fumarate ..... 35 g/ton

GUARANTEED ANALYSIS

Crude Protein (min.) 22.00%
Lysine (min.) 1.30%
Crude Fat (min.) 5.00%
Crude Fiber (max.) 3.00%
Calcium (min.) ..... 0.50% (max.) 0.75%
Phosphorus (min.) 0.55%
Salt (min.) ..... 0.25% (max.) 0.75%
Selenium (min.) ..... 0.3 ppm
Zinc (min.) ..... 175 ppm
Vitamin A (min.) .....3,250 IU/lb
Vitamin E (min.) ..... 25 IU/lb

INGREDIENTS

Ground Corn, Dehulled Soybean Meal, Lignin Sulfonate, Dried Whey, Vegetable Oil, Molasses Products, Fish Meal, Sugar, Animal Fat, Calcium Phosphate, Spray Dried Animal Blood Cells, Bentonite, Calcium Carbonate, Salt, L-Lysine, Chromium Picolinate, Zinc Methionine Complex, Natural and Artificial Flavors, Choline Chloride, dl-Methionine, Ascorbic Acid, Vitamin A Acetate, Vitamin B-12 Supplement, Vitamin D-3 Supplement, Vitamin E Supplement, Roughage Products, Niacin Supplement, Calcium Pantothenate, Biotin, Menadione Sodium Bisulfite Complex, Riboflavin Supplement, Folic Acid, Manganous Oxide, Zinc Oxide, Ferrous Sulfate, Copper Sulfate, Calcium Iodate, Sodium Selenite, Cobalt Carbonate, [Ethoxyquin, Propionic Acid, Tertiary Butyl Hydroquinone (preservatives)].

FEEDING DIRECTIONS: For control of swine dysentery, feed continuously as the sole ration on premises with a history of swine dysentery but where signs of disease have not yet occurred or following an approved treatment of the disease. For control of porcine proliferative enteropathies (ileitis), feed continuously as the sole ration for not less than 10 days. Always provide plenty of clean, fresh water.

WARNING: Withdraw medicated feed 2 days before slaughter.

CAUTION: Do not feed to animals other than swine. The effects of Tiamulin on swine reproductive performance, pregnancy and lactation have not been determined. Use as the only source of Tiamulin. Swine being treated with Tiamulin should not have access to feeds containing polyether ionophores (e.g., Monensin, Lasalocid, Narasin, Salinomycin and Semduramicin) as adverse reactions may occur. If signs of toxicity occur, discontinue use of medicated feeds.

SPECIAL CONSIDERATIONS: Protect medicated feeds from excessive temperatures. Do not store at warm temperatures, 104° F (40°C), for more than 7 days.

Manufactured By
TENNESSEE FARMERS COOPERATIVE
LAVERGNE, TENNESSEE 37086

ITEM NO. 95002

TAG CODE NO. 110316-95002

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

NET WEIGHT 50 lb (22.68 kg)/INVOICE

~JU
~LOT