

CO-OP PINNACLE LOW STARCH

For Broodmares, Stallions, Maintenance or Light to Moderate Performance Horses

GUARANTEED ANALYSIS

Crude Protein (min.)	12.00%
Lysine (min.)	0.70%
Methionine (min.)	0.25%
Crude Fat (min.)	6.00%
Crude Fiber (max.)	20.00%
Acid Detergent Fiber (max.)	27.00%
Neutral Detergent Fiber (max.)	43.00%
Dietary Starch (max.)	11.00%
Sugars (max.)	4.50%
Calcium (min.) 0.75% (max.)	1.25%
Phosphorus (min.)	0.50%
Magnesium (min.)	0.40%
Copper (min.)	60 ppm
Selenium (min.)	0.50 ppm
Zinc (min.)	150 ppm
Vitamin A (min.)	4,000 IU/lb
Vitamin D (min.)	500 IU/lb
Vitamin E (min.)	125 IU/lb

613

INGREDIENTS

Wheat Middlings, Alfalfa Meal, Soybean Hulls, Rice Mill Feed, Vegetable Oil with Tertiary Butyl Hydroquinone (a preservative), Molasses Products, Lignin Sulfonate (pelletting agent), Calcium Carbonate, Salt, Magnesium Oxide, Calcium Phosphate, Zinc Methionine Complex, Manganese Methionine Complex, Copper Lysine Complex, Cobalt Glucoheptonate, Lysine, dl-Methionine Hydroxy Analog Calcium, Calcium Hydroxide, Selenium Yeast, Artificial Flavor, Vitamin E Supplement, Ascorbic Acid, Choline Chloride, Biotin, d-Calcium Pantothenate, Vitamin A Supplement, Vitamin D-3 Supplement, Thiamine Mononitrate, Niacin Supplement, Riboflavin Supplement, Folic Acid, Vitamin B-12 Supplement, Copper Sulfate, Zinc Sulfate, Zinc Oxide, Manganous Oxide, Manganese Sulfate, Sodium Selenite, Cobalt Carbonate, Calcium Iodate, Propionic Acid (a preservative), Acetic Acid, Ammonium Hydroxide, Magnesium Aluminosilicate, Sodium Bentonite, Calcium Bentonite.

FEEDING DIRECTIONS: Co-op Pinnacle Low Starch is designed for horses with a sensitivity to high levels of starch. Follow recommended feeding rates below. Feed similar amounts at the same time each day to minimize digestive upset. Monitor the horse's body condition and adjust feeding practices to maintain an appropriate level of condition. If feeding a grain ration at a rate of 0.5 lb per 100 lb of body weight per day or more, divide total amount into 2 or 3 individual meals offered over 10 to 12 hours. Avoid abrupt changes in feed or forage type. Implement any changes to the diet over a 7- to 10-day transitional period. Changes in the amount of a grain ration should not exceed 1.0 to 1.5 lb per day. Provide free-choice access to clean, fresh water and salt at all times. Store feed in a dry area protected from insects and rodents.

FEEDING GUIDLINES:

	Feeding Rate*		
	Ponies (600 to 900 lb)	Average Horses (900 to 1200 lb)	Large Horses (1200 to 1500 lb)
Mature, Idle	3 to 5 lb	4 to 6 lb	5 to 7 lb
Mature, Light Work	3 to 5 lb	4 to 6 lb	5 to 7 lb
Mature, Moderate Work	5 to 7 lb	6 to 8 lb	7 to 9 lb
Mature, Heavy Work	7 to 9 lb	8 to 10 lb	9 to 11 lb
Broodmare, Early/Mid Gestation	3 to 5 lb	4 to 6 lb	5 to 7 lb
Yearling	5 to 7 lb	6 to 8 lb	7 to 9 lb

* Feeding rates are expressed as total lb of Pinnacle Low Starch per head per day. The feeding directions above assume horses are consuming at least 1.5% of their body weight in good-quality forage daily. Feeding amounts should be adjusted proportionately for horses with an ideal mature body weight of under 600 lb or over 1500 lb. Feed amounts may be adjusted up or down as needed to maintain ideal body condition. Horses consuming less than 0.4% of body weight (4 lb for a 1000 lb horse) should be switched to Pinnacle Balancer for a lower calorie, more concentrated source of essential nutrients.

Manufactured By
PROTRITION FEED, LLC
LAVERGNE, TENNESSEE 37086

ITEM NO. 319
TAG CODE NO. 012324-319

Seller warrants only that this feed conforms to the description on the label and is suitable for the purposes set forth when used in accordance with the directions for use on said label.

NET WEIGHT 50 lb (22.68 kg)/INVOICE