

DIFLUBENZURON	GROUP	15	INSECTICIDE
---------------	-------	----	-------------

CO-OP PASTURE BLOCK WITH CLARIFLY®

For Beef and Dairy Cattle, Sheep, Goats, Horses, Ponies, Donkeys and Mules on Pasture  
Prevents the development of house, stable, face and horn fly larvae in the manure of treated cattle and prevents the development of house and stable fly larvae in the manure of treated sheep, goats, horses, ponies, donkeys and mules.

ACTIVE INGREDIENT: Diflubenzuron\* (CAS #35367-38-5) ..... 0.02%  
OTHER INGREDIENTS: ..... 99.98%  
Total ..... 100.00%

\*Contains 90.8 mg of Diflubenzuron per lb

FEED INGREDIENTS
Cottonseed Meal, Corn Distillers Dried Grains with Solubles, Salt, Cane Molasses, Lignin Sulfonate.

\*\*Does not contain any prohibited materials. Does not contain any animal protein. This product was manufactured in a facility that does not handle or store animal protein products prohibited in ruminant feed as described in 21 CFR section 589.2000.

Use as directed.

GUARANTEED ANALYSIS
Crude Protein (min.) ..... 23.00%
Crude Fat (min.) ..... 2.00%
Crude Fiber (max.) ..... 14.00%
Acid Detergent Fiber (max.) ..... 19.00%
Neutral Detergent Fiber (max.) ..... 28.00%
Calcium (min.) ..... 0.10% (max.) ..... 0.60%
Phosphorus (min.) ..... 0.50%
Salt (min.) ..... 19.00% (max.) ..... 22.80%
Zinc (min.) ..... 40 ppm

KEEP OUT OF REACH OF CHILDREN  
CAUTION

EPA Reg. No. 89459-7-9078  
EPA Est. No. 9078-TN-4

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"><li>• Call a poison control center or doctor immediately for treatment advice.</li><li>• Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor.</li><li>• Do not give anything to an unconscious person.</li></ul>
IF ON SKIN:	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
IF IN EYES:	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
EMERGENCY ASSISTANCE: Have the product container or label with you when calling a poison control center, doctor or going for treatment. For additional information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), you may call 1-800-222-1222 24 hours per day, 7 days per week.	

ENVIRONMENTAL HAZARDS  
Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark.

See back of this label for Directions for Use, Resistance- Management, Feeding and Management Directions and Storage and Disposal.

ITEM NO. 96057  
TAG CODE NO. 022025-96057

NET WEIGHT 33 1/3 lb (15.11 kg)

Manufactured By  
PROTRITION FEED, LLC  
180 OLD NASHVILLE HIGHWAY  
LAVERGNE, TENNESSEE 37086

96057

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

## RESISTANCE-MANAGEMENT

For resistance-management, please note that Co-op Pasture Block with ClariFly® contains a Group 15/diflubenzuron insecticide. Any insect population may contain individuals naturally resistant to Co-op Pasture Block with ClariFly® and other Group 15 insecticides. The resistant individuals may dominate the insect population if these insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide/acaricide resistance, take the following steps:

- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact ProTriton Feed, LLC at 1-800-377-8380.

## FEEDING AND MANAGEMENT INSTRUCTIONS

Provide Co-op Pasture Block with ClariFly® as a free-choice source of supplementation and salt for your cattle, sheep, goats, horses, ponies, donkeys and mules. When used as directed, Co-op Pasture Block with ClariFly® will provide sufficient diflubenzuron to prevent the development of fly larvae in the manure of treated cattle, sheep, goats, horses, ponies, donkeys and mules. Diflubenzuron prevents the development of house, face, stable and horn flies in the manure of treated cattle and house and stable flies in the manure of treated sheep, goats, horses, ponies, donkeys and mules. Existing adult flies will not be affected. Start feeding this product early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall until cold weather restricts fly activity. Allow free-choice to cattle, sheep, goats horses, ponies, donkeys and mules.

Cattle should consume an average of 0.10 mg of diflubenzuron per kg of animal body weight per day to achieve control of house, face, stable and horn flies. Cattle should consume an average of 0.8 oz of Co-op Pasture Block with ClariFly® per 100 lb of animal body weight per day which is equivalent to 8 oz per day for a 1,000 lb cow.

Horses, ponies, donkeys and mules should consume an average of 0.15 mg of diflubenzuron per kg of animal body weight per day to achieve control of house and stable flies. Horses, ponies, donkeys and mules should consume an average of 1.2 oz of Co-op Pasture Block with ClariFly® per 100 lb of animal body weight per day which is equivalent to 12 oz per day for a 1,000 lb horse.

Sheep and goats should consume an average of 0.40 mg of diflubenzuron per kg of animal body weight per day to achieve control of house and stable flies. Sheep and goats should consume an average of 3.2 oz of Co-op Pasture Block with ClariFly® per 100 lb of animal body weight per day which is equivalent to 4 oz per day for a 125 lb sheep or goat.

To open, remove protective sheet and score several times with a knife to encourage initial consumption.

1. Ensure cattle, sheep, goats, horses, ponies, donkeys and mules are not starved for nutrients prior to feeding Co-op Pasture Block with ClariFly®.
2. Offer each cow, sheep, goat, horse, pony, donkey or mule 1 Co-op Pasture Block with ClariFly® when being housed individually. Offer cattle, horses, ponies, donkeys and mules 1 Co-op Pasture Block with ClariFly® for every 3 to 5 head. Offer sheep and goats 1 Co-op Pasture Block with ClariFly® for every 15 to 25 head.
3. Remove all other self-limiting nutritional supplements.
4. Locate Co-op Pasture Block with ClariFly® where cattle, sheep, goats, horses, ponies, donkeys and mules congregate (watering, loafing, shade areas).
5. Replenish Co-op Pasture Block with ClariFly® on a regular basis as needed.
6. If intake of Co-op Pasture Block with ClariFly® is below 0.8 oz per 100 lb animal body weight per day for cattle or 1.2 oz per 100 lb animal body weight per day for horses, ponies, donkeys and mules or 3.2 oz per 100 lb animal body weight per day for sheep and goats, increase the number of feeding locations and/or change locations to areas more frequented by cattle, sheep, goats, horses, ponies, donkeys and mules.
7. If intake of Co-op Pasture Block with ClariFly® is above 0.8 oz per 100 lb animal body weight per day for cattle or 1.2 oz per 100 lb animal body weight per day for horses, ponies, donkeys and mules or 3.2 oz per 100 lb animal body weight per day for sheep and goats, reduce the number of feeding locations and/or change locations to areas less frequented by cattle, sheep, goats, horses, ponies, donkeys and mules.
8. Introduction of this product after adult fly infestation is established may require treatment of cattle, sheep, goats, horses, ponies, donkeys and mules with supplemental sprays to reduce the adult fly population coming from outside sources.

**STORAGE AND DISPOSAL:** Do not contaminate water, food or feed by storage or disposal.

**Storage:** Store in a cool, dry place.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Completely empty wrapper. Dispose of wrapper in a sanitary landfill or by other procedures approved by state and local authorities.

For information or in an emergency call 1-800-222-1222.

Made in USA